

Quintessential brane cosmology

K.E. Kunze¹

*Département de Physique Théorique, Université de Genève
24 Quai Ernest Ansermet, CH-1211 Genève 4, Switzerland*

and

M.A. Vázquez-Mozo^{2,3}

*Niels Bohr Institute, Blegdamsvej 17,
DK-2100 Copenhagen Ø, Denmark*

We study a class of braneworlds where the cosmological evolution arises as the result of the movement of a three-brane in a five-dimensional static dilatonic bulk, with and without reflection symmetry. The resulting four-dimensional Friedmann equation includes a term which, for a certain range of the parameters, effectively works as a quintessence component, producing an acceleration of the universe at late times. Using current observations and bounds derived from big-bang nucleosynthesis we estimate the parameters that characterize the model.

September 2001

¹E-mail: kunze@amorgos.unige.ch

²E-mail: vazquez@nbi.dk.

³Address after January 1st, 2002: Theory Division, CERN, CH-1211 Geneva 23, Switzerland.

1 Introduction and motivation

One of the most interesting experimental results in cosmology in the last years has been the measure of the acceleration of the universe from the observation of high-redshift supernovae [1]. The combination of this result with the measure of the cosmic microwave background (CMB) anisotropies [2] favors a universe whose energy density is very close to the critical one, $\Omega_{\text{total}} \simeq 1$, and with a large dark energy component, $\Omega_{\text{dark}} \simeq 0.7$. This realization that the energy content of the universe might be dominated by some kind of dark energy has revived the interest in cosmological models where the late time expansion of the universe is governed by some kind of vacuum energy, either in the form of a nonvanishing cosmological constant or a quintessence field (see [3, 4] for comprehensive reviews).

Recently, braneworld scenarios have emerged as a very appealing alternative to standard dimensional reduction in higher-dimensional theories. The general idea is the following. The observable universe is a submanifold embedded in a higher-dimensional space-time. By some physical mechanism matter fields are confined to this submanifold. Although gravity can propagate into the bulk, the Kaluza-Klein excitations of the four-dimensional graviton can be decoupled at low energies either by considering finite but “large” extra dimensions [5] or a negative cosmological constant in the bulk [6]. Therefore, at energies below the mass gap of these Kaluza-Klein modes, physics on the brane is well described by the matter degrees of freedom confined to the brane and standard gravity in four dimensions. One of the most interesting features of these models is that the energy scale at which the five-dimensional effects are important can be relatively low, even of the order of the TeV, and therefore in principle accessible to future high-energy experiments.

It has been realized that the braneworld paradigm in its cosmological application leads to corrections to the four-dimensional Einstein equations on the brane [7, 8, 9]. These departures from General Relativity, and their imprints in braneworld cosmological models, might be used to test the scenario against cosmological observations. In the context of Friedmann-Robertson-Walker cosmologies, the braneworld hypothesis results in the presence of two extra terms on the right-hand side of the four-dimensional Friedmann equation: a dark-radiation contribution and a quadratic term in the energy density that couples directly to the five-dimensional Planck scale. While the first one might have any effect at the present stage of the history of the universe, the second one is expected to be important only in the

very early universe where the energy density was very large.

Quintessence models have also been studied in the context of brane cosmologies [10, 11]. With a few exceptions [12], in most of the models considered a four-dimensional decaying cosmological constant is implemented by introducing an additional scalar field in four dimensions. By a suitable choice of potential its contribution to the energy density becomes dominant only at present.

In this paper, we study the possibility within a braneworld scenario that an effective four-dimensional quintessence is induced by a dilaton field propagating in the bulk. In our model the cosmological evolution in four dimensions reflects the movement of the braneworld in a static dilatonic bulk space-time in which $\mathbb{Z}_2$ -reflection symmetry around the location of the brane can be broken. We find that the bulk dilaton induces a term in the effective four-dimensional Einstein equations that acts as a decaying cosmological constant. Hence, in this sense quintessence occurs naturally in this class of models. For universes filled with matter and radiation, and for a certain range of the parameters defining the model, this vacuum energy actually tracks the matter or radiation energy density and dominates at late times, driving an accelerating expansion of the universe. Using constraints from big-bang nucleosynthesis and the present value of Newton's constant allows us to constrain the parameters of our class of models. In particular, we find a lower bound of 10^5 GeV for the five-dimensional Planck mass.

In the solutions showing quintessencelike behavior the universe first undergoes decelerated expansion and then at late times enters into a regime of accelerated expansion due to the effects of the bulk dilaton. In this case, the observed isotropy of the universe has to be assumed. However, as will be seen, the class of models under study also allows for inflationary solutions at early times. In this type of solution the universe starts in a state of accelerated expansion and then later undergoes decelerated expansion. The problem in this kind of models, however, is not only the lack of acceleration in the expansion at late times, but the fact that in this asymptotic regime the decelerated expansion is dominated by the ρ^2 term in the Friedmann equation.

Much of the recent interest in brane cosmologies was triggered by Hořava-Witten type models [13] where the brane is confined to lie on the fixed points of an $\mathbb{S}^1/\mathbb{Z}_2$ orbifold. Thus usually a $\mathbb{Z}_2$ symmetry in the bulk direction is assumed. However here we will consider more

general situations in which the bulk space-time is not symmetric under reflection around the position of the brane. In our context, this situation describes braneworlds that emerge as domain walls separating two regions of space-time with different values of the cosmological constant. As we will see, this asymmetry leaves its imprints in the cosmological dynamics on the brane at late times, for example through corrections to the four-dimensional Newton's constant depending on the difference between the cosmological constants on the two sides of the brane.

The plan of the paper is as follows. In Section 2 we study the dynamics of moving braneworlds in the background of five-dimensional Einstein-Liouville solutions. Section 3 will be devoted to the study of the cosmological dynamics of moving brane universes with $\mathbb{Z}_2$ -reflection symmetry, whereas in Section 4 we will analyze brane models where this reflection symmetry is broken. In Section 5 we will proceed to study the phenomenology of these models by using both current observational data and bounds derived from big-bang nucleosynthesis. Finally, in Section 6, we will summarize our results.

2 Moving branes in a static bulk geometry

In Ref. [14] a broad class of braneworld cosmologies coupled to a dilaton field in the bulk was studied by using a solution generating algorithm that enables the construction of five-dimensional Einstein-Liouville cosmologies (cf. [18]). Assuming $\mathbb{Z}_2$ -reflection symmetry around the fixed location of the brane, it was found that the solutions studied enjoy a new version of the self-tuning mechanism of [19], in which the total cosmological constant in four-dimensions is canceled independently of the value of the bulk cosmological constant and the energy density on the brane. It was also found there that inflation in Friedmann-Robertson-Walker and Bianchi-I models belonging to the family of solutions studied cannot be achieved for any value of the amplitude and slope of the bulk Liouville potential.

Here we are going to use these bulk solutions in a different physical setup. Our aim is the construction of realistic four-dimensional brane cosmologies in which the cosmological expansion accelerates at late times. For this, even if we restrict the bulk solutions to those studied in [14], we will have to relax some of the conditions imposed on the braneworlds constructed there. In particular, here we will not consider the braneworld to have a fixed location on the bulk, but we will take into account the degree of freedom associated with

its collective motion in the five-dimensional space-time. In addition, we will also consider situations in which $\mathbb{Z}_2$ -reflection symmetry is broken (cf. [15, 16, 17]). As we will see in the following, the features of the resulting brane cosmologies are quite different from the ones studied in [14].

2.1 Description of the model

We will consider solutions to the five-dimensional bulk action

$$S_{5D} = \frac{1}{\kappa_5^2} \int d^5x \sqrt{-\mathcal{G}} \left[\frac{1}{2} \mathcal{R} - \frac{1}{2} (\partial\phi)^2 - \Lambda e^{-\frac{2}{3}k\phi} \right] + \int_Y d^4x \sqrt{-g} \left[\frac{1}{\kappa_5^2} K^\pm - \lambda(\phi) + e^{4b\phi} L(e^{2b\phi} g_{\mu\nu})_{\text{matter}} \right]. \quad (2.1)$$

Here we are assuming that the five-dimensional Planck scale κ_5 determines also the scale of the five-dimensional scalar field, which is dimensionless here. The only other independent energy scales entering into the action (2.1) are provided by Λ , which has dimensions of (mass)², and the energy scale of the four-dimensional dynamics. In the following, we will consider that the matter on the brane is described by a perfect fluid with barotropic equation of state $p = w\rho$ (with $|w| \leq 1$), so the matter Lagrangian is written in terms of the energy-density as $L_{\text{matter}} = -\rho$. In [14], solutions to the equations of motion in the bulk derived from (2.1) were constructed in terms of the functions $Q(x)$ and $h_{\mu\nu}(x)$ in which the metric takes the form⁴.

$$ds_{5D}^2 = \frac{1}{\xi^2} e^{\frac{4k}{\sqrt{k^2+6}}Q(x)} r^{\frac{2}{3}(k^2-3)} dr^2 + e^{-\frac{2k}{\sqrt{k^2+6}}Q(x)} r^2 h_{\mu\nu}(x) dx^\mu dx^\nu \quad (2.2)$$

with the scalar field given by

$$\phi(x, r) = \frac{6}{\sqrt{k^2+6}} Q(x) + k \log r. \quad (2.3)$$

The metric function $h_{\mu\nu}$ is a solution to the four-dimensional Einstein equations coupled to a scalar field, $R_{\mu\nu} = \partial_\mu \psi \partial_\nu \psi$ with $\psi(x) = \sqrt{6}Q(x)$. The amplitude of the Liouville potential as a function of k and ξ has the following form

$$\Lambda = \frac{1}{2}(k^2 - 12)\xi^2. \quad (2.4)$$

⁴These solutions are related to the ones studied in [14] by a k -dependent coordinate redefinition $r^2 = \exp(a_2\chi)$ and a shift in the dilaton field.

The parameter ξ has dimensions of mass and determines the scale of the cosmological constant.

Here we will study the motion of a braneworld in a bulk geometry described by the line element (2.2), possibly with different values of ξ on the two sides of the brane. In order to simplify our analysis, we consider $Q(x) \equiv 0$ in Eq. (2.3) and the bulk space-time to be static. This implies in particular that $h_{\mu\nu}$ is a static Ricci-flat four-dimensional metric. Hence the metric takes the form

$$ds_{5D}^2 = \frac{1}{\xi^2} r^{\frac{2}{3}(k^2-3)} dr^2 + r^2 h_{\mu\nu} dx^\mu dx^\nu, \quad (2.5)$$

while the dilaton is

$$\phi = k \log r. \quad (2.6)$$

In addition, the four-dimensional metric is given by

$$h_{\mu\nu} dx^\mu dx^\nu = -dt^2 + h_{ij}(x) dx^i dx^j. \quad (2.7)$$

Let us then consider a single braneworld embedded in the five-dimensional space-time described by Eq. (2.5) and moving along the trajectory

$$t = t(\tau), \quad r = R(\tau), \quad x, y, z = \text{constant}, \quad (2.8)$$

where τ is the proper time. For a five-dimensional static metric of the form $ds^2 = \mathcal{G}_{55} dr^2 + \mathcal{G}_{00} dt^2 + \mathcal{G}_{ij} dx^i dx^j$, the nonvanishing components of the 5-velocity $\mathcal{V}^A$ and the vector normal to the brane n^A can be computed using the relations $\mathcal{V}_A \mathcal{V}^A = -1$, $\mathcal{V}_A n^A = 0$, with the result

$$\begin{aligned} \mathcal{V}^0 &= \sqrt{\frac{1 + \mathcal{G}_{55} \dot{R}^2}{-\mathcal{G}_{00}}}, & \mathcal{V}^5 &= \dot{R} \\ n^0 &= \dot{R} \sqrt{\frac{\mathcal{G}_{55}}{-\mathcal{G}_{00}}}, & n^5 &= \sqrt{\frac{1 + \mathcal{G}_{55} \dot{R}^2}{\mathcal{G}_{55}}}. \end{aligned} \quad (2.9)$$

Here and in the rest of the paper the dots indicate differentiation with respect to proper time τ .

On the other hand, taking an orthonormal frame $\{\mathcal{V}_M, n_M, e_M^{(i)}\}$, the components of the extrinsic curvature can be easily computed using the existence of the timelike Killing vector of the static bulk space-time (cf. [20, 21]):

$$K_\tau^\tau = \frac{1}{\dot{R} \sqrt{-\mathcal{G}_{00} \mathcal{G}_{55}}} \frac{d}{d\tau} \left[\sqrt{-\mathcal{G}_{00}} \sqrt{1 + \mathcal{G}_{55} \dot{R}^2} \right], \quad (2.10)$$

$$K_j^i = \frac{1}{2} n^M \mathcal{G}^{ik} \partial_M \mathcal{G}_{kj}. \quad (2.11)$$

Note that $K_{\tau\tau}$ is given by the orthogonal component of the proper acceleration of the brane.

Looking at Eqs. (2.5) and (2.7), we see that, when expressed in proper time, the induced metric on the brane takes the form

$$ds_{4D}^2 = -d\tau^2 + R^2(\tau)h_{ij}(x)dx^i dx^j. \quad (2.12)$$

Since the metric (2.7) has to be a static vacuum solution to Einstein equations, in the following we will focus on the simplest possibility where $h_{ij} = \delta_{ij}$. In this case the line element (2.12) describes a flat Friedmann-Robertson-Walker universe.

2.2 Junction conditions and dynamical equations

As explained above, we will consider that the constant ξ determining the amplitude of the Liouville potential takes in general different values $\xi_{\pm}$ on the two sides of the brane (a related proposal has been studied in [16]). This means that the bulk geometry is described by the line elements

$$\begin{aligned} ds_{5D}^2 \Big|_+ &= \frac{1}{\xi_+^2} r^{\frac{2}{3}(k^2-3)} dr^2 + r^2[-dt^2 + \delta_{ij}dx^i dx^j], & r > R(\tau), \\ ds_{5D}^2 \Big|_- &= \frac{1}{\xi_-^2} r^{\frac{2}{3}(k^2-3)} dr^2 + r^2[-N(t)^2 dt^2 + \delta_{ij}dx^i dx^j], & r < R(\tau). \end{aligned} \quad (2.13)$$

Following [16] we have introduced a shift function $N(t)$ in order to ensure the continuity of the metric across the brane. Notice that the bulk field equations are satisfied since, given the static character of the metric, the shift function can be locally reabsorbed in a redefinition of time. A simple calculation shows that the metric is continuous across $r = R(\tau)$ by taking

$$N(t)^2 = \left(\frac{\xi_-}{\xi_+}\right)^2 \frac{\xi_+^2 + R^{\frac{2}{3}(k^2-3)} \dot{R}^2}{\xi_-^2 + R^{\frac{2}{3}(k^2-3)} \dot{R}^2}. \quad (2.14)$$

On the other hand, the dilaton is continuous across the location of the brane.

The Israel junction conditions, together with the matching for the dilaton across the brane, will determine the dynamics of the scale factor $R(\tau)$. As for the metric junction equations we find

$$[K_{\mu\nu}] \equiv K_{\mu\nu}^+ - K_{\mu\nu}^- = \frac{\kappa_5^2}{3} [-\lambda(\phi) + g^{\alpha\beta}\tau_{\alpha\beta}] g_{\mu\nu} - \kappa_5^2 \tau_{\mu\nu}, \quad (2.15)$$

where $g_{\mu\nu}$ is the induced metric, $\tau_{\mu\nu}$ is the energy-momentum tensor on the brane and $\lambda(\phi)$ is the brane tension. Indices are raised and lowered using $g_{\mu\nu}$. Since we are assuming

that the matter on the brane is a perfect fluid the energy-momentum tensor is given by $\tau_\nu^\mu = \text{diag}(-\rho, p, p, p)$. Using the barotropic equation of state $p = w\rho$ we find from the matching condition for K_τ^τ ,

$$\begin{aligned} & \left[\left(\frac{\dot{R}}{R} \right)^2 + \xi_+^2 R^{-\frac{2}{3}k^2} \right]^{-\frac{1}{2}} \left[\frac{\ddot{R}}{R} + \frac{k^2}{3} \left(\frac{\dot{R}}{R} \right)^2 + \xi_+^2 R^{-\frac{2}{3}k^2} \right] \\ & - \left[\left(\frac{\dot{R}}{R} \right)^2 + \xi_-^2 R^{-\frac{2}{3}k^2} \right]^{-\frac{1}{2}} \left[\frac{\ddot{R}}{R} + \frac{k^2}{3} \left(\frac{\dot{R}}{R} \right)^2 + \xi_-^2 R^{-\frac{2}{3}k^2} \right] = -\frac{\kappa_5^2}{3} \lambda(\phi) + \kappa_5^2 \left(w + \frac{2}{3} \right) \rho \end{aligned} \quad (2.16)$$

while from the jump of K_j^i at $r = R(\tau)$ we get

$$\left[\left(\frac{\dot{R}}{R} \right)^2 + \xi_+^2 R^{-\frac{2}{3}k^2} \right]^{\frac{1}{2}} - \left[\left(\frac{\dot{R}}{R} \right)^2 + \xi_-^2 R^{-\frac{2}{3}k^2} \right]^{\frac{1}{2}} = -\frac{\kappa_5^2}{3} [\lambda(\phi) + \rho]. \quad (2.17)$$

In addition to this, we have also to impose the matching condition for the dilaton derivative across the brane [21],

$$[n^A \partial_A \phi] = \kappa_5^2 [\lambda'(\phi) - b g^{\mu\nu} \tau_{\mu\nu}] \quad (2.18)$$

which, upon substitution, reads

$$k \left[\left(\frac{\dot{R}}{R} \right)^2 + \xi_+^2 R^{-\frac{2}{3}k^2} \right]^{\frac{1}{2}} - k \left[\left(\frac{\dot{R}}{R} \right)^2 + \xi_-^2 R^{-\frac{2}{3}k^2} \right]^{\frac{1}{2}} = \kappa_5^2 [\lambda'(\phi) - (3w - 1)b\rho]. \quad (2.19)$$

The particular case in which the bulk space-time has $\mathbb{Z}_2$ -reflection symmetry across the position of the brane is recovered by setting $\xi_+ = \xi_- \equiv \xi$ and taking the outward normals on the two sides of the brane with opposite signs. In Eqs. (2.16)-(2.19), this amounts to replacing the minus sign between the two terms on the left hand side of these equations by a plus sign.

Equations (2.16)-(2.19) derived from the junction conditions determine the dynamics of the four-dimensional scale factor $R(\tau)$ in terms of the brane tension $\lambda(\phi)$ and the energy density ρ . Consistency of Eqs. (2.17) and (2.19) implies the following differential equation relating the brane tension with the energy density:

$$\lambda'(\phi) + \frac{k}{3} \lambda(\phi) = \left[(3w - 1)b - \frac{k}{3} \right] \rho. \quad (2.20)$$

It is remarkable that, because of the junction condition for the dilaton, the matter energy density on the brane actually mixes with the brane tension.

It is important to stress that any solution of Eq. (2.17) will automatically satisfy the second order differential equation derived from Eq. (2.16), provided the brane tension $\lambda(\phi)$ and the density ρ satisfy (2.20) and (2.22). Thus, as it is the case in standard cosmology, the dynamics of the brane universe is solely determined by the first-order differential equation (2.17).

The analysis of the case $k = 0$ requires some care. Since for vanishing k the dilaton field exactly vanishes for the family of solutions studied, the dilaton matching equation (2.19) is not imposed and the only dynamical equation is Eq. (2.17) with constant brane tension $\lambda(\phi) = \lambda_0$. In this case, therefore, there is no mixing between the vacuum energy and the matter on the brane.

2.3 Energy balance on the brane

In order to get the form of the brane tension $\lambda(\phi)$ from Eq. (2.20) it would be necessary to know how the energy density depends on the scale factor (or equivalently, for $k \neq 0$, on the dilaton field). The conservation equations for the energy-momentum on the brane can be derived from the Codazzi equation. Actually, as explained in [8], in the presence of a dilaton field there is an interchange of energy-momentum between the brane and the bulk. This implies the existence of a nonvanishing right-hand side in the conservation equation

$$\dot{\rho} + 3\frac{\dot{R}}{R}(\rho + p) = -(3p - \rho) b \dot{R} \frac{d\phi}{dR}. \quad (2.21)$$

We see that the source term in the continuity equation results from the conformal coupling between the matter-energy density on the brane and the bulk dilaton, measured by b .

Using the equation of state of the fluid and the expression of the dilaton field in terms of the scale factor, Eq. (2.6), we find the following differential equation for $\rho(R)$:

$$R \frac{d\rho}{dR} + m\rho = 0, \quad \text{with} \quad m \equiv 3(w + 1) + (3w - 1) k b. \quad (2.22)$$

This implies that the energy density depends on the scale factor and the dilaton field as

$$\rho(R) = \rho_0 R^{-m} \quad \Longrightarrow \quad \rho(\phi) = \rho_0 e^{-\frac{m}{k}\phi}, \quad (2.23)$$

where the second equation is only valid for $k \neq 0$. By comparing these expressions with the ones encountered in ordinary cosmology, we find that the usual evolution of the energy density with the scale factor is corrected by a term that depends on the conformal coupling b between the dilaton and the matter fields. Equation (2.23) can be physically interpreted by considering that, for an observer on the brane the source term in Eq. (2.25) leads to an effective barotropic index w_{eff} governing the evolution of the energy density with the scale factor $\rho(R) \sim R^{-3(w_{\text{eff}}+1)}$, where

$$w_{\text{eff}} = w + \left(w - \frac{1}{3}\right) kb. \quad (2.24)$$

In Section 5, we will see that for phenomenologically viable models $k^2 \lesssim 10^{-2}$. Since it is natural to expect the coupling b to be of the same order as k , the difference between w and w_{eff} would be also of order 10^{-2} . There are, however, two particular instances in which the correction to w in Eq. (2.24) vanishes for any value of b . The first one is when $k = 0$, which corresponds to switching the dilaton off, so the right hand side of Eq. (2.21) is zero and the usual continuity equation for $\rho(R)$ is recovered. The second one corresponds to the case in which we consider a radiation fluid on the brane, $w = \frac{1}{3}$, where again the right-hand side of Eq. (2.21) vanishes. This is a consequence of the classical conformal invariance of radiation, which is therefore insensitive to the conformal coupling to the dilaton. As a result, when the matter on the brane is dominated by a gas of massless particles, the temperature evolves with the scale factor in the same way as in radiation-dominated standard cosmology, $T(R) \sim R^{-1}$.

By plugging the expression for $\rho(\phi)$ into the right hand side of Eq. (2.20) we find a differential equation for the brane tension. When $k^2 \neq 0, 3m$, the solution is given by

$$\lambda(\phi) = \lambda_0 e^{-\frac{k}{3}\phi} - \left[1 + \frac{9(w+1)}{k^2 - 3m}\right] \rho_0 e^{-\frac{m}{k}\phi}. \quad (2.25)$$

where λ_0 is an integration constant. In the particular case when $k^2 = 3m$ the solution takes the form

$$\lambda(\phi)_{k^2=3m} = \left[\lambda_0 - \frac{3}{k}(w+1)\rho_0\phi\right] e^{-\frac{k}{3}\phi}. \quad (2.26)$$

Finally, if $k = 0$ the brane tension is constant and does not mix with the matter-energy density,

$$\lambda(\phi)_{k=0} = \lambda_0. \quad (2.27)$$

In the following, unless stated otherwise, we will only study the case when $k^2 \neq 3m$.

As pointed out above, it is interesting that the expressions (2.25) and (2.26) for the brane tension depend on the matter energy density through the constant ρ_0 . This mixing between matter and vacuum energies on the brane can be seen to be dilaton-mediated, in the sense that it is produced by the junction conditions of the dilaton on the brane, which couples to both the brane vacuum energy (through its dependence on ϕ) and the matter fields in the action (2.1). Only when $k = 0$ is the dilaton switched off and there is no mixing between the matter and the vacuum energy on the brane.

3 Moving braneworlds in a $\mathbb{Z}_2$ -symmetric bulk space-time

In this section, we are going to analyze the cosmological dynamics of a four-dimensional braneworld moving on a bulk space-time with $\mathbb{Z}_2$ -reflection symmetry around the location of the brane. This case is of some interest, since it represents a deformation of the cosmological models studied in [14] by taking into account the degrees of freedom associated with the movement of the brane in the bulk.

The dynamical equations for the scale factor are obtained from Eq. (2.17) by taking $\xi_+ = \xi_- \equiv \xi$ and reversing the minus sign on the left hand side. This results in

$$\left(\frac{\dot{R}}{R}\right)^2 = -\xi^2 R^{-\frac{2}{3}k^2} + \frac{\kappa_5^4}{36}(\lambda + \rho)^2. \quad (3.1)$$

As explained in the previous section, there is a mixing between the brane tension $\lambda(\phi)$ and the energy density due to the mediating dilaton. This implies that the expression of the tension as a function of the dilaton field (2.25) contains a term depending on ρ . If we collect all the ρ -dependent terms in Eq. (3.1) together we can write the Friedmann-like equation ($k^2 \neq 3m$)

$$\begin{aligned} \left(\frac{\dot{R}}{R}\right)^2 &= \frac{1}{36}(\kappa_5^4 \lambda_0^2 - 36\xi^2)R^{-\frac{2}{3}k^2} + \frac{\kappa_5^4}{18} \left(\pm \lambda_0 R^{-\frac{k^2}{3}}\right) \left|\frac{9(w+1)}{k^2 - 3m}\right| \rho(R) \\ &+ \frac{\kappa_5^4}{36} \left[\frac{9(w+1)}{k^2 - 3m}\right]^2 \rho(R)^2, \end{aligned} \quad (3.2)$$

where the $\pm$ signs correspond to $k^2 < 3m$ and $k^2 > 3m$, respectively. Let us also remark here

that in order for the (time-dependent) Newton constant to be positive, we have to restrict $\lambda_0 > 0$ when $k^2 < 3m$ and $\lambda_0 < 0$ if $k^2 > 3m$.

When $w > -1$ we can transform Eq. (3.2) into the usual Friedmann equation in brane cosmology (cf. [22, 23]) by defining the effective energy density in four dimensions through the rescaling,

$$\rho(R)_{\text{eff}} = \left| \frac{9(w+1)}{k^2 - 3m} \right| \rho(R), \quad (3.3)$$

which amounts to a redefinition of the (positive) integration constant ρ_0 in Eq. (2.23). In physical terms, this means that the energy density measured by a four-dimensional observer by looking at the rate of expansion differs from the “bare” value that appears in the action (2.2) by a positive multiplicative constant. This effective energy density satisfies the same continuity equation as $\rho(R)$ and scales in the same way with the scale factor. We will identify it with the energy density driving the cosmological evolution of our brane universe.

In order to simplify the notation in the following, we will drop the subscript “eff” and denote the effective four-dimensional energy density by $\rho(R)$. Doing so we finally arrive at the Friedmann equation on the brane,

$$\left(\frac{\dot{R}}{R} \right)^2 = \frac{8\pi}{3} G_N(R) \rho_Q(R) + \frac{8\pi}{3} G_N(R) \rho(R) + \frac{\kappa_5^4}{36} \rho(R)^2. \quad (3.4)$$

We see that, in addition to the usual terms linear and quadratic in $\rho(R)$, there is also a “quintessence” energy density given by

$$\rho_Q(R) = \frac{\kappa_5^4 \lambda_0^2 - 36\xi^2}{2\kappa_5^4 |\lambda_0|} R^{-\frac{k^2}{3}}. \quad (3.5)$$

As we did in Eq. (2.24), we can define the effective barotropic index associated with the quintessence term that determines the scaling of ρ_Q with the scale factor as

$$w_Q = \frac{k^2}{9} - 1. \quad (3.6)$$

Thus, for $k^2 \leq 6$, it satisfies $-1 \leq w_Q \leq -\frac{1}{3}$ and represents a real quintessence contribution to the Friedmann equation. In addition to this, the four-dimensional effective gravitational coupling in our braneworlds depends on the scale factor according to

$$G_N(R) = \frac{\kappa_5^4 |\lambda_0|}{48\pi} R^{-\frac{k^2}{3}}. \quad (3.7)$$

In the case $w = -1$ and $k^2 \neq 0$, the Friedmann equation is retrieved directly from Eq. (3.2), with the result

$$\left(\frac{\dot{R}}{R}\right)^2 = \frac{1}{36}(\kappa_5^4 \lambda_0^2 - 36\xi^2)R^{-\frac{2}{3}k^2} \equiv \frac{8\pi}{3}G_N(R)\rho_Q(R), \quad (3.8)$$

so the dependence on the matter energy density disappears and the dynamics of the universe is controlled only by the induced “quintessence” energy density. When $w = -1$ and $k = 0$, there is an explicit dependence on ρ_0 and the corresponding Friedmann equation is given by (3.8) with the replacement $\lambda_0 \rightarrow \lambda_0 + \rho_0$.

In order to study the solutions of Eq. (3.4), it is convenient to rewrite it as the equation of motion of a particle moving with zero total energy in a potential,

$$\dot{R}^2 + V(R) = 0, \quad (3.9)$$

where

$$V(R) = \xi^2 R^{-\frac{2}{3}(k^2-3)} - \frac{\kappa_5^4}{36} \left[|\lambda_0| R^{-\frac{1}{3}(k^2-3)} + \rho_0 R^{-(m-1)} \right]^2. \quad (3.10)$$

Inflationary solutions on the brane are those with $\dot{R} > 0$, $\ddot{R} > 0$. Using the equations of motion (3.9), we find that the conditions for inflation reduce to

$$V'(R) < 0, \quad (3.11)$$

where the prime denotes differentiation with respect to R .

By looking at Eq. (3.10), we see that if $m > 1$, the brane energy density tends to stop inflation, whereas the bulk and brane vacuum energies compete with each other. When $k^2 < 3$, the brane vacuum energy contributes to accelerate the expansion, whereas the bulk cosmological constant tends to decelerate it. If $k^2 > 3$ the situation is reversed. In the following we will study more explicitly the main features of the cosmological solutions obtained in the cases when the matter on the brane is dominated by either vacuum energy ($w = -1$) or a gas of ultrarelativistic particles ($w = \frac{1}{3}$).

3.1 Vacuum branes ($w = -1$)

Let us first consider the case in which the matter on the brane contributes to the energy-momentum tensor as a vacuum energy with the equation of state $p = -\rho$. When $k \neq 0$, we

find from Eq. (3.8) that all dependence on ρ_0 disappears from the Friedmann equation and the potential $V(R)$ simplifies to

$$V(R) = \left(\xi^2 - \frac{\kappa_5^4}{36} \lambda_0^2 \right) R^{-\frac{2}{3}(k^2-3)}. \quad (3.12)$$

In order to have solutions to Eq. (3.9) we must demand $\kappa_5^4 \lambda_0^2 \geq 36\xi^2$. When the inequality is saturated ($\kappa_5^4 \lambda_0^2 = 36\xi^2$) the brane remains at a fixed position $R(\tau) = \text{constant}$ and we retrieve Minkowski space-time on the brane for any value of k . The brane tension is given by

$$\lambda(\phi) = \pm \frac{6\xi}{\kappa_5^4} e^{-\frac{k}{3}\phi} - \rho_0 e^{4b\phi} \quad (3.13)$$

and the energy-momentum tensor on the brane is $\tau_{\mu\nu} = -\rho_0 e^{4b\phi} \eta_{\mu\nu}$ (cf. [14]). Actually, these are the self-tuning solutions of Ref. [19] which preserve four-dimensional Poincaré invariance for any value of the cosmological constant in the bulk.

If, on the other hand, $\kappa_5^4 \lambda_0^2 > 36\xi^2$ we find that $V'(R) < 0$ for $k^2 < 3$ and hence inflation takes place. By solving explicitly the equations of motion (3.9) with the potential (3.12), we find that when $k^2 \neq 0$, the scale factor scales with the proper time as

$$R(\tau) \sim \tau^{\frac{3}{k^2}}. \quad (3.14)$$

Thus, for $0 < k^2 < 3$, the four-dimensional universe undergoes power-law inflation.

On the other hand, if $k = 0$, the brane tension is constant $\lambda(\phi) = \lambda_0$ and solutions to Eq. (3.9) will exist provided $\kappa_5^4 (\lambda_0 + \rho_0)^2 \geq 36\xi^2$. When this inequality is saturated, we find the static Randall-Sundrum model [6]. Otherwise, the four-dimensional metric is not static and we have de Sitter inflation,

$$R(\tau) \sim \exp \left(\tau \sqrt{\frac{1}{3} \Lambda_{\text{eff}}} \right), \quad (3.15)$$

i.e., in this case the expansion is driven by an effective four-dimensional cosmological constant $\Lambda_{\text{eff}} = \frac{\kappa_5^4}{12} (\lambda_0 + \rho_0)^2 - 3\xi^2 > 0$.

3.2 Radiation branes ($w = \frac{1}{3}$)

The second physically interesting case is the one in which the matter on the branes is a perfect fluid with equation of state $p = \frac{1}{3}\rho$. The first interesting thing to notice in this case

is that $m = 4$ and all the equations become independent of the parameter b measuring the coupling between the dilaton and the matter on the brane. This is a consequence of the classical conformal invariance of a radiation perfect fluid. Now the potential $V(R)$ is given by

$$V(R) = \xi^2 R^{-\frac{2}{3}(k^2-3)} - \frac{\kappa_5^4}{36} \left[|\lambda_0| R^{-\frac{1}{3}(k^2-3)} + \rho_0 R^{-3} \right]^2 \quad (3.16)$$

with $\lambda_0 > 0$ for $k^2 < 12$ and $\lambda_0 < 0$ if $k^2 > 12$.

In order to extract the qualitative behavior of the scale factor, we can study the asymptotic limits for large and small scale factors. When $k^2 < 12$, we find that

$$V(R) \sim \begin{cases} \left(\xi^2 - \frac{\kappa_5^4}{36} \lambda_0^2 \right) R^{-\frac{2}{3}(k^2-3)}, & R \gg \left(\frac{\rho_0}{|\lambda_0|} \right)^{\frac{3}{12-k^2}}, \\ -\frac{\kappa_5^4}{36} \rho_0^2 R^{-6}, & R \rightarrow 0. \end{cases} \quad (3.17)$$

“Large” universes will result only if we demand $\kappa_5^4 \lambda_0^2 > 36 \xi^2$. In this case, we observe that for large values of R the evolution of the scale factor is identical to the one found for vacuum branes. Thus, for $k = 0$, the universe undergoes de Sitter inflation while for $0 < k^2 < 3$ inflation occurs with the scale factor scaling as a power of the proper time. When $k^2 = 3$, we find coasting behavior of the scale factor and for $3 < k^2 < 12$ the expansion of the four-dimensional universe is decelerated.

On the other hand, when $R \rightarrow 0$ the evolution is always decelerated with $R(\tau) \sim \tau^{\frac{1}{4}}$ for all $0 < k^2 < 12$. This is the expected evolution of the scale factor of a radiation universe whose dynamics is dominated by the ρ^2 term in Eq. (3.4). It is interesting to note that the evolution of the universe in this regime is completely dominated by the matter energy density, while for large R the expansion rate is determined by a competition between the bulk cosmological constant (encoded in ξ) and the brane vacuum energy as measured by λ_0 .

Altogether, we find that for a radiation filled braneworld in a bulk dilaton potential with $k^2 < 12$, the cosmological evolution in four dimensions begins with a ρ^2 -dominated phase. After a period of decelerated expansion resulting in a decrease of the density, the universe will enter a regime in which expansion is driven by the term proportional to ρ in the Friedmann equation. This “standard” expansion phase is terminated at some time in which the matter energy density decreases below the contribution of the energy density from the quintessence term. At this moment, the universe will start its accelerated expansion. To illustrate the point, in Fig. 1 we have plotted the potential $V(R)$ and the scale factor for a particular

model in this class of solutions. In Section 5 we will study this type of brane universe as quintessence models for the observed acceleration of the universe.




Figure 1: Potential $V(R)$ (left) and numerical solution to the scale factor $R(\tau)$ (right) for a model with $k = 1$, $\xi = 0.1$, $\lambda_0 = 1$ and $\rho_0 = 1$. The five-dimensional Planck scale is taken $\kappa_5 = 1$.

Incidentally, when $k^2 < 12$ and $\kappa_5^4 \lambda_0^2 > 36\xi^2$ ($\lambda_0 > 0$) the universe starts at a big-bang singularity at $\tau = 0$ and recollapses after reaching a maximum size of $R_{\max} = [\kappa_5^4 \rho_0 / (6\xi + \kappa_5^4 |\lambda_0|)]^{\frac{3}{12-k^2}}$.

Let us analyze the case when $k^2 > 12$ which corresponds to a positive bulk cosmological constant. Now the asymptotic behavior of the potential is

$$V(R) \sim \begin{cases} -\frac{\kappa_5^4}{36} \rho_0^2 R^{-6}, & R \gg \left(\frac{\rho_0}{|\lambda_0|} \right)^{\frac{3}{12-k^2}}, \\ \left(\xi^2 - \frac{\kappa_5^4}{36} \lambda_0^2 \right) R^{-\frac{2}{3}(k^2-3)}, & R \rightarrow 0 \end{cases} \quad (3.18)$$

and the situation is reversed with respect to the case analyzed before. For large values of R , the potential is always negative with $V'(R) > 0$ and the decelerated evolution is controlled by the matter energy density through the ρ^2 term in the Friedmann equation (3.4). If again we demand $\kappa_5^4 \lambda_0^2 < 36\xi^2$ ($\lambda_0 < 0$), we find that the universe does not have a big-bang singularity, but it starts at a finite size to undergo an inflationary expansion in which the scale factor grows as a power law. This regime is followed by a “graceful exit” phase where the universe enters a decelerated expansion which asymptotically leads to the ρ^2 term dominance. An example of this kind of solutions is presented in Fig. 2. In the left panel we have plotted the

potential $V(R)$ and on the right one a numerical solution of Eq. (3.9) is shown. The total amount of inflation is just determined by the ratio between the size of the universe at the end of inflation $V'(R_f) = 0$ and the original size of the universe $V(R_i) = 0$.




Figure 2: Profile of the potential $V(R)$ (left) and numerical solution of the scale factor $R(\tau)$ (right) for $k = 4$, $\xi = 1$, $\lambda_0 = -1$ and $\rho_0 = 3$. We have also taken $\kappa_5 = 1$.

If instead we have $\kappa_5^4 \lambda_0^2 > 36\xi^2$, the universe starts at $R = 0$ and experiences a decelerated evolution to reach asymptotically the ρ^2 -dominated regime. This difference in the behavior when $R \rightarrow 0$ can be understood by noticing that in the previous case the dynamics of the universe at small values of R is dominated by a large negative vacuum energy that prevents the formation of the singularity.

Perhaps the most surprising feature of the solutions with $k^2 > 12$ is the fact that the late-time (large R) regime is dominated by the ρ^2 term in (3.4). This might seem counter-intuitive, since it is usually argued that this term should only be important in the very early universe where the energy density of the universe is very large. The key element to understand the large- R behavior of these solutions is to realize that in our model the four-dimensional gravitational coupling evolves with time as $R^{-\frac{k^2}{3}}$, whereas the ρ^2 term couples directly to the five-dimensional Planck scale, which is time independent. Thus if k is large enough, both the “quintessence” term and the term linear in the energy density become subleading with respect to the ρ^2 contribution when the scale factor grows. Therefore, the cosmological evolution is asymptotically dominated by the bulk term in the Friedmann equation.

Before finishing with the analysis of radiation branes, we study, for the sake of completeness, the marginal case $k^2 = 12$ where the bulk dilaton potential vanishes. As explained at the end of Section 2, when $k^2 = 3m \equiv 12$ the brane tension is given by Eq. (2.26), which contains a logarithmic dependence on R . Then $V(R)$ is given by

$$V(R)_{k^2=12} = \left[\xi^2 - \frac{\kappa_5^4}{36} (\lambda_0 + \rho_0 - 4\rho_0 \log R)^2 \right] R^{-6}. \quad (3.19)$$

This implies that for both large and small R , the dynamics of the scale factor is controlled by the logarithmic term in the potential. The profile of the potential presents then two allowed regions for a zero-energy particle, for large and small R , separated by a barrier.

4 Cosmologies on the brane in bulks with broken $\mathbb{Z}_2$ -reflection symmetry

As we discussed in Section 2, braneworlds can be also embedded in a bulk metric of the form (2.2) without $\mathbb{Z}_2$ -reflection symmetry by considering different values of the energy scale ξ at the two sides of the brane location. In this case, the evolution equations for the scale factor are obtained by solving Eq. (2.17) for the Hubble parameter with the result

$$\left(\frac{\dot{R}}{R} \right)^2 = -\frac{\xi_+^2 + \xi_-^2}{2} R^{-\frac{2}{3}k^2} + \frac{\kappa_5^4}{36} (\lambda + \rho)^2 + \frac{9}{4\kappa_5^4} (\xi_+^2 - \xi_-^2)^2 (\lambda + \rho)^{-2} R^{-\frac{4}{3}k^2}. \quad (4.1)$$

Comparing with Eq. (3.1), we notice the presence of an extra term which scales with a negative power of the brane energy density (cf. [17]). The expressions for $\rho(R)$ and $\lambda(\phi)$ are the same as for the symmetric case. Proceeding as in the previous section, we reduce the problem of integrating the scale factor to finding the trajectory of a classical nonrelativistic particle with zero energy in the potential,

$$\begin{aligned} V(R) &= \frac{\xi_+^2 + \xi_-^2}{2} R^{-\frac{2}{3}(k^2-3)} - \frac{\kappa_5^4}{36} \left[|\lambda_0| R^{-\frac{1}{3}(k^2-3)} + \rho_0 R^{-(m-1)} \right]^2 \\ &\quad - \frac{9}{4\kappa_5^4} (\xi_+^2 - \xi_-^2)^2 R^{-\frac{4}{3}(k^2-3)} \left[|\lambda_0| R^{-\frac{1}{3}(k^2-3)} + \rho_0 R^{-(m-1)} \right]^{-2}, \end{aligned} \quad (4.2)$$

where, as above, $\rho_0 R^{-m}$ represents the rescaled effective energy density on the brane. We notice that the last extra term in the potential associated with the breaking of reflection symmetry is always negative and therefore its contribution tends to increase the value of

$H^2 \equiv \left(\frac{\dot{R}}{R}\right)^2$. Moreover, it can be easily checked that this new term is bounded by the second one on the right-hand side of Eq. (4.2) for all values of R provided $|\lambda_0| \geq \frac{3}{\kappa_5^2} \sqrt{|\xi_+^2 - \xi_-^2|}$. As a consequence, when this bound is satisfied the dynamics of the brane universe is never dominated by the extra term in Eq. (4.1) and the cosmological evolution is qualitatively similar to the one found in the previous section.

In order to show the qualitative differences with respect to the symmetric case studied in the previous section, we have to evaluate the relevance of the extra term in Eq. (4.1). Here we will focus on the solutions with phenomenological applications, therefore, for reasons that will be explained in Section 5, we restrict our attention to cases with $k^2 < 3m$. If $|\lambda_0| \geq \frac{3}{\kappa_5^2} \sqrt{|\xi_+^2 - \xi_-^2|}$, the extra term is negligible at small R compared with the dominating ρ^2 term. In general, however, accelerated expansion takes place for large values of R . In this latter case, the effective four-dimensional vacuum energy depends on the scale of $\mathbb{Z}_2$ -symmetry breaking, $(\xi_+^2 - \xi_-^2)^2$,

$$V(R) \sim \begin{cases} \left[\frac{\xi_+^2 + \xi_-^2}{2} - \frac{\kappa_5^4}{36} \lambda_0^2 - \frac{9(\xi_+^2 - \xi_-^2)^2}{4\kappa_5^4 \lambda_0^2} \right] R^{-\frac{2}{3}(k^2-3)}, & R \gg \left(\frac{\rho_0}{|\lambda_0|} \right)^{\frac{3}{3m-k^2}}, \\ -\frac{\kappa_5^2}{36} \rho_0^2 R^{-2(m-1)}, & R \rightarrow 0. \end{cases} \quad (4.3)$$

Taking $\kappa_5^4 \lambda_0^2 [18(\xi_+^2 + \xi_-^2) - \kappa_5^4 \lambda_0^2] < 81(\xi_+^2 - \xi_-^2)^2$ it is easy to check that the universe recollapses within finite proper time without passing through a phase of accelerated expansion. Therefore, these solutions are not very interesting from a phenomenological point of view. On the other hand, if $\kappa_5^4 \lambda_0^2 [18(\xi_+^2 + \xi_-^2) - \kappa_5^4 \lambda_0^2] > 81(\xi_+^2 - \xi_-^2)^2$, the universe starts again in a ρ^2 -dominated phase that, when ρ decreases, will be followed by a standard expansion dominated by the term linear in ρ in Eq. (4.1). Eventually, however, the (positive) quintessence energy density will dominate and the universe enters a regime of accelerated expansion.

Actually, when R is large enough and higher powers of ρ can be neglected [which includes the limit $R \gg (\rho_0/|\lambda_0|)^{\frac{3}{3m-k^2}}$] the dynamics of the universe can be derived from the approximate Friedmann equation,

$$\left(\frac{\dot{R}}{R}\right)^2 \approx \frac{8\pi}{3} G_N(R) \rho_Q(R) + \frac{8\pi}{3} G_N(R) \rho(R), \quad (4.4)$$

where the new Newton constant is given by

$$G_N(R) \approx \frac{\kappa_5^4 |\lambda_0|}{48\pi} \left[1 - \frac{81}{2\kappa_5^8 \lambda_0^4} (\xi_+^2 - \xi_-^2)^2 \right] R^{-\frac{k^2}{3}} \quad (4.5)$$

and the quintessence component of the energy density is

$$\rho_Q(R) \approx \frac{1}{2\kappa_5^4|\lambda_0|} \left[\kappa_5^4\lambda_0^2 - 18(\xi_+^2 + \xi_-^2) + \frac{162}{\kappa_5^4\lambda_0^2} \left(1 - 9\frac{\xi_+^2 + \xi_-^2}{\kappa_5^4\lambda_0^2} \right) (\xi_+^2 - \xi_-^2)^2 \right] R^{-\frac{k^2}{3}}. \quad (4.6)$$

For consistency, only terms linear in $\frac{1}{\kappa_5^4\lambda_0^2}(\xi_+ - \xi_-)^2$ are retained upon substitution in Eq. (4.4). We see that both Newton's constant and the quintessence energy density reduce to expressions (3.5) and (3.7) for $\xi_+ = \xi_-$.

Let us consider now the case in which the difference between the cosmological constant at the two sides of the brane is “large”, i.e. $\kappa_5^2|\lambda_0| < 3\sqrt{|\xi_+^2 - \xi_-^2|}$. Now the last term in Eq. (4.1) will dominate for a certain range of values of R . Thus, the ρ^2 -dominated epoch at small R will be followed by a regime where the expansion is driven by the term proportional to $(\lambda + \rho)^{-2}$. This will happen for the following range of values of the scale factor:

$$\left(\frac{\kappa_5^2\rho_0}{3\sqrt{|\xi_+^2 - \xi_-^2|}} \right)^{\frac{3}{3m-k^2}} \lesssim R \lesssim \left(\frac{\rho_0}{|\lambda_0|} \right)^{\frac{3}{3m-k^2}}. \quad (4.7)$$

For larger values of R , the universe will enter a quintessence-dominated regime.

To summarize, we have seen that when the difference between the values of ξ at both sides of the brane is small (in the sense explained above), the cosmological dynamics of the universe does not show a radically different behavior with respect to the symmetric models. Actually, for large values of R the effect of the breaking of the reflection symmetry shows in a correction of both Newton's constant and ρ_Q , although their scalings with the scale factor are not affected. On the other hand, for larger values of $|\xi_+ - \xi_-|$, the main difference is the existence of an intermediate phase interpolating between the ρ^2 regime and the “standard” expansion dominated by the linear term in the Friedmann equation, where the dynamics of the universe is controlled by the last term in Eq. (4.2). At large values of R , however, the dynamics can still be described by the approximate Friedmann equation (4.4).

5 Phenomenology of quintessential brane cosmologies

After having investigated their mathematical properties, in the following we will study phenomenological aspects of the quintessential brane cosmological models presented in Section 3, where the acceleration of the universe at late times is induced by the rolling of the braneworld in a static bulk geometry. We will focus our attention on the $\mathbb{Z}_2$ -reflection symmetric models.

We saw that whenever $\kappa_5^4 \lambda_0^2 > 36\xi^2$ and $k^2 < 3m$ (with $m > 0$), solutions describing a flat Friedmann-Robertson-Walker universe can be found which, after going through radiation/matter-dominated epochs, undergo accelerated expansion at late times. For these types of solutions, the energy content of the very early universe is dominated by radiation and the expansion is controlled by the ρ^2 term in Eq. (3.4). When the density is low enough, a phase of “standard” expansion starts where the dynamics of the universe is governed by the term linear in ρ in the Friedmann equation. It is during this phase that nucleosynthesis has to take place, as well as the decoupling of matter from radiation. Finally, some time after the universe enters in a matter dominated epoch, the quintessence term starts dominating the dynamics and the universe begins its accelerated expansion. This acceleration is driven by the bulk scalar field through the term ρ_Q in the Friedmann equation, which, as we will see below, plays the role of effective quintessence on the brane. In addition to this, the effective Newton constant in our models, Eq. (3.7), evolves with time through its dependence on the scale factor.

We normalize the scale factor on the brane so it takes the value $R_0 = 1$ at the present time and we use the temperature of the radiation to label the different epochs in the evolution of the universe,

$$T(R) = \frac{2.73 \text{ K}}{R}. \quad (5.1)$$

From our results in Sec. 2.3 we saw that the energy density on the brane scales with the temperature as

$$\rho(T) = \rho_0 \left(\frac{T}{2.73 \text{ K}} \right)^{3(1+w_{\text{eff}})}, \quad (5.2)$$

where w_{eff} is give by Eq. (2.24). As discussed above, for the particular case of radiation $w_{\text{eff}} = w = \frac{1}{3}$, and we recover Boltzmann’s law, $\rho(T) \simeq T^4$. However, when $w = 0$ the effective barotropic index is nonvanishing. Nonetheless, in the following we will denominate “matter” to any perfect fluid with $w_{\text{eff}} = 0$, whose energy therefore scales with the temperature as $\rho(T) \simeq T^3$, as seen by a brane observer. Assuming by naturality that $b \simeq k$, we find that $w_{\text{eff}} - w \simeq k^2$. As we will see later, this number is of order $\lesssim 10^{-2}$ so the difference between the “microscopic” barotropic index and the effective one (which governs the cosmological evolution) is very small.

To study the evolution of the universe in more quantitative terms, it is useful to introduce three parameters, $\Theta_Q(T)$, $\Theta_\rho(T)$, and $\Theta_{\rho^2}(T)$, which measure the contribution to the expan-

sion of the three terms on the right hand side of the Friedmann equation (3.4), normalized to the Hubble parameter,

$$\begin{aligned}\Theta_Q(T) &= \frac{8\pi}{3H^2}G_N(T)\rho_Q(T), \\ \Theta_\rho(T) &= \frac{8\pi}{3H^2}G_N(T)\rho(T), \\ \Theta_{\rho^2}(T) &= \frac{\kappa_5^4\rho(T)^2}{36H^2},\end{aligned}\tag{5.3}$$

where $\rho_Q(T)$ is given by Eq.(3.5). In this way the Friedmann equation is written as

$$1 = \Theta_Q(T) + \Theta_\rho(T) + \Theta_{\rho^2}(T).\tag{5.4}$$

The crossover between the ρ^2 -dominated epoch and the standard regime will take place at a temperature T_c at which $\Theta_{\rho^2}(T_c) \simeq \Theta_\rho(T_c)$. By using the definitions (5.3) we find

$$\frac{T_c}{2.73\text{ K}} \simeq \left(\frac{2|\lambda_0|}{\rho_0}\right)^{\frac{3}{12-k^2}}.\tag{5.5}$$

On the other hand the standard expansion of the universe will be terminated at a temperature $T_Q < T_c$ in which $\Theta_\rho(T_Q) \simeq \Theta_Q(T_Q)$. Since this has to occur after the universe has entered a matter-dominated era, we find

$$\frac{T_Q}{2.73\text{ K}} \simeq \left(\frac{\kappa_5^4\lambda_0^2 - 36\xi^2}{2\kappa_5^4|\lambda_0|\rho_0}\right)^{\frac{3}{9-k^2}}.\tag{5.6}$$

As a consequence, the period of standard expansion can be very large if the net cosmological constant on the brane, $\kappa_5^4\lambda_0^2 - 36\xi^2$, is small enough.

In order to illustrate the discussion above, in Fig. 3 the evolution of the Hubble parameter is shown for some “realistic” values of the parameters in a cold dark matter model. We can see clearly several regimes. At high temperatures ($T \gtrsim 10^{12}\text{ K}$), the expansion is dominated by the ρ^2 term in the Friedmann equation. This is followed by a long period of standard expansion for $10^{12}\text{ K} \gtrsim T \gtrsim 10\text{ K}$. In this regime, radiation ($m = 4$) dominates the expansion for temperatures higher than $T_{\text{eq}} \simeq 10^4\text{ K}$, whereas for lower temperatures the energy density of the universe is dominated by matter ($m = 3$). Finally, when $T \lesssim 10\text{ K}$, the quintessence component takes over and the universe starts its accelerated expansion.




Figure 3: Evolution of the Hubble parameter (GeV) vs. temperature (K) for a realistic quintessence model. The values of the parameters are: $\rho_0 = 0.43\rho_{Q,0} = 2.43 \times 10^{-47} \text{ GeV}^4$, $G_{N,0} = 6.71 \times 10^{-39} \text{ GeV}^{-2}$, $T_{\text{eq}} = 10^4 \text{ K}$, $k^2 = 0.01$ and $\kappa_5 = 10^{-8} \text{ GeV}^{-\frac{3}{2}}$.

5.1 Bounds from big-bang nucleosynthesis

The abundances with which light elements are synthesized in the early universe are extremely sensitive to changes in the expansion rate of the universe and/or in the fundamental constants [24, 25]. Thus, primordial nucleosynthesis provides a good way to bound the values of the constants and scales that characterize a given cosmological model. In the case at hand, the effective gravitational coupling varies with time and from Eq. (3.7) we can see that this change is determined solely by k , namely

$$\frac{\dot{G}_N}{G_N} = -\frac{k^2}{3}H. \quad (5.7)$$

In order to explain the primordial abundances of ^4He and other light elements, however, Newton's constant at the time of nucleosynthesis cannot be very different from the present value. Actually, its time variation has to satisfy [26]

$$\left(\frac{\dot{G}_N}{G_N}\right)_0 \lesssim \pm 0.01 h^{-1} H_0. \quad (5.8)$$

Therefore, we find that the parameter k that determines the slope of the potential of the bulk dilaton has to be rather small,

$$k^2 \lesssim 0.03 h^{-1}, \quad (5.9)$$

which, from Eq. (3.6), implies $w_Q + 1 \lesssim 0.003 h^{-1}$ (cf. [27]).

A second condition to be met by a cosmological model in order to account for the primordial abundances of the light elements is that the energy density of the universe at the time of nucleosynthesis has to be dominated by radiation. In particular, the dark energy contribution to the density parameter satisfies the following bound [28] (see also [29] and references therein):

$$\Omega(T_{\text{BBN}})_{\text{dark}} < 0.045 \quad (\text{at } 2\sigma \text{ confidence level}), \quad (5.10)$$

where $T_{\text{BBN}} \simeq 10^{10} \text{K}$. In our model, the dark energy component corresponds to the contribution of ρ_Q . Thus, we define

$$\Omega_Q(T) = \frac{\rho_Q(T)}{\rho_Q(T) + \rho(T)}, \quad (5.11)$$

where $\rho(T)$ is the matter/radiation energy density.

From Eq. (5.11), and assuming a cold dark matter scenario, we can obtain an expression of $\Omega_Q(T_{\text{BBN}})$ in terms of quantities at the present time as

$$\Omega_Q(T_{\text{BBN}}) = \left[1 + \frac{1 - \Omega_Q(T_0)}{\Omega_Q(T_0)} \left(\frac{T_{\text{eq}}}{T_0} \right)^3 \left(\frac{T_{\text{BBN}}}{T_{\text{eq}}} \right)^4 \left(\frac{T_{\text{BBN}}}{T_0} \right)^{-\frac{k^2}{3}} \right]^{-1}, \quad (5.12)$$

where $T_{\text{eq}} \simeq 10^4 \text{K}$ is the temperature at which the density of radiation and matter are equal, and $T_0 = 2.73 \text{K}$ is the temperature of the CMB at the present time. Since the measurement of CMB anisotropies and the supernova data indicate that $\Omega_Q(T_0) \simeq 0.7$, we find that for models satisfying the bound (5.9) the contribution of the dark energy component at the time of nucleosynthesis is

$$\Omega_Q(T_{\text{BBN}}) \simeq 10^{-35}, \quad (5.13)$$

which falls well inside the nucleosynthesis bound (5.10).

5.2 Bounds from observations

We have seen that by satisfying the bound (5.9), our quintessential brane cosmologies comply with the constraints imposed by big-bang nucleosynthesis. It is interesting to notice, however, that none of the two bounds considered in the previous section involves any other parameter of the model but the numerical constant k , or equivalently the quintessence effective barotropic index w_Q . In particular we do not get any information as to the size of the dimensionful parameters λ_0 and ξ or the five-dimensional Planck scale κ_5 .

In order to extract bounds for these scales, we have to compare the values of ρ_Q and the present value of the four-dimensional Newton constant with the observed values. From Eq. (3.7) we get the value of the Newton's constant by setting $R_0 = 1$. Using $G_{N,0} = 6.71 \times 10^{-39} \text{ GeV}^{-2}$, we find

$$\kappa_5^4 |\lambda_0| = 6\kappa_4^2 \simeq 10^{-36} \text{ GeV}^{-2}. \quad (5.14)$$

The second observational result we can use to estimate the scales of the model is the value of the dark energy component of the density parameter, $\Omega(T_0)_{\text{dark}} \simeq 0.7$. From Eqs. (5.11) and (3.5), we find that

$$\kappa_5^4 \lambda_0^2 - 36\xi^2 \simeq 4.67\kappa_5^4 |\lambda_0| \rho_0, \quad (5.15)$$

where ρ_0 is the matter density of the universe at the present time. Using Eq. (5.14) and the value $\rho_0 \simeq 2.43 \times 10^{-47} h^2 \text{ GeV}^4$ [24] we arrive at

$$\kappa_5^4 \lambda_0^2 - 36\xi^2 \simeq 10^{-82} \text{ GeV}^2. \quad (5.16)$$

Incidentally, using Eqs. (5.15) and (5.6) we find that $T_Q \simeq 3.63 \text{ K}$.

Equations (5.14) and (5.16) are the only two estimates that we can get just using “in-brane” information. In order to go forward and get some estimate for the five-dimensional Planck scale κ_5 or the brane tension λ_0 , we need some knowledge of the fundamental energy scale in the higher-dimensional theory. The ρ^2 term in the Friedmann equation (3.4) provides a window to the bulk dynamics, since it couples directly to the five-dimensional Planck scale κ_5 . Therefore, it would be very interesting to have some information about the transition between the ρ^2 -dominated epoch and the standard expansion. Even if we do not have a concrete value for the temperature at which this transition takes places, we can still get a

bound for λ_0 by going back to Eq. (5.5). In order to preserve big-bang nucleosynthesis, the ρ^2 -component in the Friedmann equation has to be negligible at T_{BBN} [30]. Imposing $T_c \gg 10^{10}$ K, we are led to

$$|\lambda_0| \gg 10^{40} \rho_0 \simeq 10^{-7} \text{ GeV}^4. \quad (5.17)$$

Using Eq. (5.14) the following bound for the five-dimensional Planck mass is found,

$$M_5 \equiv \kappa_5^{-\frac{2}{3}} \gg 10^5 \text{ GeV}, \quad (5.18)$$

and consequently $\xi \gg 10^{-22} \text{ GeV}$. Notice that this bound for the five-dimensional Planck mass is stronger than the one obtained in the quintessence scenario of Ref. [11], although it is again weaker than the one obtained from the Randall-Sundrum model [6].

6 Conclusions

In the present paper, we have studied the bulk solutions constructed in Ref. [14] in a different physical setup. Instead of being at a fixed location, the braneworlds move in a five-dimensional dilatonic bulk space time with and without $\mathbb{Z}_2$ -reflection symmetry. As a result, and contrary to the case analyzed in Ref. [14], the cosmological dynamics is not driven by a time-dependent background but by the movement of the braneworld in an otherwise static bulk space-time. The resulting class of brane cosmologies admits, depending on the choice of parameters, either inflationary behavior at early times and decelerated expansion at late times or a decelerated expansion at early times which evolves into a quintessencelike behavior at late, i.e., present, times.

The realization of quintessence in these models seems to be particularly interesting since it does not rely on the introduction of an additional scalar field on the brane. It is in fact driven by the bulk dilaton. As the brane moves along in the static bulk, it sweeps out sections with different values for the bulk dilaton and, from the point of view of an observer living on the brane, this is interpreted as a time-dependent cosmological constant. For a suitable choice of parameters, this evolution is such that the energy density of this effective quintessence only becomes important at the present time. However, in addition to the induced time-dependent cosmological constant the bulk dilaton also causes a time-dependent gravitational coupling. Using observational constraints on the variation of Newton's constant and the present value of the cosmological constant gives bounds on the parameter of the brane cosmologies at hand.

The fact that $\kappa_5^4 \lambda_0^2 - 36\xi^2 \ll \kappa_5^4 \lambda_0^2, \xi^2$ seems to indicate that the nonvanishing value of the four-dimensional vacuum energy is the result of a slightly broken symmetry. Furthermore, we find a lower limit for the five-dimensional Planck mass of 10^5 GeV.

Here we are taking a phenomenological approach and therefore we do not attempt to explain these values for the scales of the model from first principles. However, it would be very interesting to investigate whether the hierarchy of scales found here can be obtained in a natural way by embedding these five-dimensional models in a string/M-theory setup.

We have shown that for a certain range of parameters, inflation takes place in the early universe. However, this phase is followed, after a graceful exit period, by a decelerated Friedmann-Robertson-Walker universe in which the expansion is driven by ρ^2 . This fact makes this particular set of solutions not very promising as phenomenological models of the universe [31].

We have also considered brane cosmologies with broken $\mathbb{Z}_2$ -reflection symmetry and found that for a slight symmetry breaking the evolution follows qualitatively that of the symmetric case. In this case, for large values of the scale factor the evolution is approximately described by a Friedmann equation where both Newton's constant and the quintessence energy density receive corrections depending on the scale of $\mathbb{Z}_2$ -symmetry breaking.

One of the interesting properties of the braneworlds investigated in [14] is the existence of an extension of the mechanism of self-tuning [19] in which the matching conditions at the brane location force the cancellation of all vacuum energy terms on the right-hand side of the Einstein equations in four dimensions. However, from our analysis in Section 3 we find that the physical mechanism enforcing this self-tuning is the condition that the braneworld is at a fixed position in the bulk. Indeed, from Eq. (3.1) we conclude that the condition $\dot{R} = 0$ for any R is equivalent to consider a vacuum brane ($w = -1$) with zero net cosmological constant in four dimensions, $\kappa_5^4 \lambda_0^2 = 36\xi^2$. From this point of view, at least in the case of a static bulk geometry, the self-tuning can be seen as resulting from the tuning of the velocity of the brane in the bulk to zero⁵. However, this is not necessarily a fine-tuning, since the condition $\dot{R} = 0$ can be automatically implemented if the braneworld is confined to lie on the fixed points of an $\mathbb{S}^1/\mathbb{Z}_2$ orbifold, as it is the case in Hořava-Witten type models [13].

⁵ This result extends to the cases with $\mathbb{Z}_2$ -reflection symmetry breaking, where again a nonmoving brane is only achieved by considering a vacuum brane with $\kappa_5^4 \lambda_0^2 = 9(\xi_+ \pm \xi_-)^2$.

Acknowledgments

It is a pleasure to thank Alex Feinstein for interesting discussions and a most enjoyable collaboration in Ref. [14]. K.E.K. has been supported by the Swiss National Science Foundation and Spanish Science Ministry Grant 1/CICYT 00172.310-0018-12205/2000. The work of M.A.V.-M. has been supported by EU Network “Discrete Random Geometry” Grant HPRN-CT-1999-00161, ESF Network no. 82 on “Geometry and Disorder”, Spanish Science Ministry Grant AEN99-0315, and University of the Basque Country Grants UPV 063.310-EB187/98 and UPV 172.310-G02/99. K.E.K. thanks the Niels Bohr Institute for hospitality. M.A.V.-M. thankfully acknowledges the hospitality of the Theory Division at CERN and the Cosmology Group at the Physics Department of the University of Geneva during the completion of this work.

References

- [1] S. Perlmutter *et al.*, *Nature* **391** (1998) 51 ([astro-ph/9712212](#)); *Astrophys. J.* **517** (1999) 565 ([astro-ph/9812133](#)).
A.G. Riess *et al.*, *Astron. J.* **116** (1998) 1009 ([astro-ph/9805201](#)).
- [2] P. de Bernardis *et al.* *Nature* **404** (2000) 955 ([astro-ph/0004404](#));
A. Balbi *et al.*, *Astrophys. J.* **545** (2000) L1 ([astro-ph/0005124](#)).
- [3] S. Weinberg, *Rev. Mod. Phys.* **61** (1989) 1; *The cosmological constant problems*, talk at Dark Matter 2000, [hep-ph/0005265](#);
N. Straumann, *On the mystery of the cosmic vacuum energy density*, talk at DARK 2000, [astro-ph/0009386](#);
A. Vilenkin, *Cosmological constant problems and their solutions*, talk at The Dark Universe: Matter, Energy, and Gravity, [hep-th/0106083](#).
- [4] P. Binetruy, *Int. J. Theor. Phys.* **39** (2000) 1859 ([hep-ph/0005037](#)).
- [5] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett.* **B429** (1998) 263 ([hep-th/9803315](#));
I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett.* **B436** (1998) 257 ([hep-th/9804398](#)).
- [6] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83** (1999) 3370 ([hep-th/9905221](#)); *Phys. Rev. Lett.* **83** (1999) 4690 ([hep-th/9906064](#)).

- [7] T. Shiromizu, K. Maeda and M. Sasaki, *Phys. Rev.* **D62** (2000) 024012 (gr-qc/9910076);
- [8] K. Maeda and D. Wands, *Phys. Rev.* **D62** (2000) 124009 (hep-th/0008188).
- [9] A. Mennim and R.A. Battye, *Class. Quantum Grav.* **18** (2001) 2171 (hep-th/0008192).
- [10] P.F. González-Díaz, *Phys. Lett.* **B481** (2000) 353 (hep-th/0002033);
 G. Huey and J.E. Lidsey, *Phys. Lett.* **B514** (2001) 215 (astro-ph/0104006);
 M. Gasperini, *Phys. Rev.* **D64** (2001) 043510 (gr-qc/0105082);
 A.S. Majumdar, *Phys. Rev.* **D64** (2001) 083503 (astro-ph/0105518);
 B. Chen and F.-L. Lin, *Holographic cosmic quintessence on dilatonic brane world*, hep-th/0106054;
 A. Albrecht, C.P. Burgess, F. Ravndal and C. Skordis, *Natural quintessence and large extra dimensions*, astro-ph/0107573;
 Y.S. Myung, *Quintessence and brane world scenarios*, hep-th/0107065;
 M. Gasperini, F. Piazza and G. Veneziano, *Quintessence as a run-away dilaton*, gr-qc/0108016;
 Y.S. Myung, *Mod. Phys. Lett.* **A16** (2001) 1963 (hep-th/0108092).
- [11] K. Maeda, *Phys. Rev.* **D64** (2001) 123525 (astro-ph/0012313);
 S. Mizuno and K. Maeda, *Phys. Rev.* **D64** (2001) 123521 (hep-ph/0108012).
- [12] P. Brax and A.C. Davis, *Phys. Lett.* **B497** (2001) 289 (hep-th/0011045); *J. High Energy Phys.* **05** (2001) 007 (hep-th/0104023);
 D. Langlois and M. Rodríguez-Martínez, *Phys. Rev.* **D64** (2001) 123507 (hep-th/0106245);
 C. Charmousis, *Dilaton space-times with a Liouville potential*, hep-th/0107126.
- [13] P. Hořava and E. Witten, *Nucl. Phys.* **B 460** (1996) 506 (hep-th/9510209); *ibid.* **B 475** (1996) 94 (hep-th/9603142).
- [14] A. Feinstein, K.E. Kunze and M.A. Vázquez-Mozo, *Phys. Rev.* **D64** (2001) 084015 (hep-th/0105182).
- [15] P. Kraus, *J. High Energy Phys.* **12** (1999) 011 (hep-th/9910149);
 A. Kehagias and E. Kiritsis, *J. High Energy Phys.* **11** (1999) 22 (hep-th/9910174);
 H. Stoica, S.-H. Henry Tye and I. Wasserman, *Phys. Lett.* **B482** (2000) 205 (hep-th/0004126);
 B. Carter and J.-P. Uzan, *Nucl. Phys.* **B606** (2001) 45 (gr-qc/0101010);

- R.A. Battye, B. Carter, A. Mennim and J.-P. Uzan, *Phys. Rev.* **D64** (2001) 124007 (hep-th/0105091).
- [16] N. Deruelle and T. Doležal, *Phys. Rev.* **D62** (2000) 103502 .
- [17] A.-C. Davis, S. C. Davis, W.B. Perkins and I.R. Vernon, *Phys. Lett.* **B504** (2001) 254 (hep-ph/0008132); *Phys. Rev.* **D63** (2001) 083518 (hep-ph/0012223).
- [18] J.E. Lidsey, *Class. Quantum Grav.* **17** (2000) L39 (gr-qc/9911066).
- [19] S. Kachru, M. Schulz and E. Silverstein, *Phys. Rev.* **D62** (2000) 045021 (hep-th/0001206).
- [20] M. Visser, *Lorentzian wormholes: From Einstein to Hawking*, AIP, New York, 1995; C. Barceló and M. Visser, *Nucl. Phys.* **B584** (2000) 415 (hep-th/0004022).
- [21] H.A. Chamblin and H.S. Reall, *Nucl. Phys.* **B562** (1999) 133 .
- [22] P. Binetruy, C. Deffayet and D. Langlois, *Nucl. Phys.* **B565** (2000) 269 (hep-th/9905012).
- [23] P. Binetruy, C. Deffayet, U. Ellwanger and D. Langlois, *Phys. Lett.* **B477** (2000) 285 (hep-th/9910219).
- [24] J.A. Peacock, *Cosmological Physics*, Cambridge 1999.
- [25] K.A. Olive, G. Steigman and T.P. Walker, *Phys. Rep.* **333** (2000) 389 (astro-ph/9905320)
- [26] E.W. Kolb, M.J. Perry and T.P. Walker, *Phys. Rev.* **D33** (1986) 869 ;
F. S. Accetta, L.M. Krauss and P. Romanelli, *Phys. Lett.* **B248** (1990) 146.
- [27] P.S. Corasaniti and E.J. Copeland, *Constraining the quintessence equation of state with $SnIa$ data and CMB peaks*, astro-ph/0107378.
- [28] R. Bean, S.H. Hansen and A. Melchiorri, *Phys. Rev.* **D64** (2001) 103508 (astro-ph/0104162).
- [29] P.G. Ferreira and M. Joyce, *Phys. Rev. Lett.* **79** (1997) 4740 (astro-ph/9707286);
Phys. Rev. **D58** (1998) 023503 (astro-ph/9711102).
- [30] S.M. Carroll and M. Kaplinghat, *Testing the Friedmann equation: the expansion of the universe during big-bang nucleosynthesis*, astro-ph/0108002.

- [31] R. Maartens, D. Wands, B.A. Bassett and I. Heard, *Phys. Rev.* **D62** (2000) 041301 (hep-ph/9912464)
- E.J. Copeland, A.R. Liddle and J.E. Lidsey, *Phys. Rev.* **D64** (2001) 023509 (astro-ph/0006421)
- H. Kodama, A. Ishibashi and O. Seto, *Phys. Rev.* **D62** (2000) 064022 (hep-th/0004160);
- C. van de Bruck, M. Dorca, R.H. Brandenberger and A. Lukas, *Phys. Rev.* **D62** (2000) 123515 (hep-th/0005032);
- D. Langlois, *Phys. Rev.* **D62** (2000) 126012 (hep-th/0005025); *Phys. Rev. Lett.* **86** (2001) 2212 (hep-th/0010063);
- C. Gordon and R. Maartens, *Phys. Rev.* **D63** (2001) 044022 (hep-th/0009010);
- N. Deruelle, T. Doležal and J. Katz, *Phys. Rev.* **D63** (2001) 083513 (hep-th/0010215);
- T. Boehm, R. Durrer and C. van de Bruck, *Phys. Rev.* **D64** (2001) 063504 (hep-th/0102144);
- S.M. Carroll and L. Mersini, *Phys. Rev.* **D64** (2001) 124008 (hep-th/0105007);
- V. Sahni, M. Sami and T. Souradeep, *Relic gravity waves from brane world inflation* gr-qc/0105121;
- V. Bozza, M. Gasperini and G. Veneziano, *Localization of scalar fluctuations in a dilatonic brane world scenario*, hep-th/0106019;
- P. Brax, C. van de Bruck and A.C. Davis, *J. High Energy Phys.* **10** (2001) 026 (hep-th/0108215)